

Beam-beam effects: modelling, measurements and correction strategy on the luminosity calibration measurements at the Large Hadron Collider experiments

Tatiana Pieloni and Joanna Wańczyk (EPFL)

X. Buffat (CERN), A. Dabrowski (CERN), W. Kozanecki (IRFU-CEA), D. Stickland (Princeton U.), R. Tomas Garcia (CERN), Y. Wu (EPFL), A. Babaev (TPU), C. Tambasco, A. Mehta, O. Karacheban (CERN), G. Pasztor, (Eotvos, U) T. Barklow (SLAC), I. Kralik (IEF), J. Wenninger, LHC OP crews

This work is supported by the Swiss Accelerator Research and Technology Institute (CHART)

**HB2023 workshop
CERN, 10th October**

Luminosity Basics

$$N_{events} = L \times \sigma_{event}$$

$\mu_{vis} = \epsilon * \mu =$ mean number of interactions per Bunch
crossing seen by detector

$$L = \frac{\mu_{vis} n_b f_r}{\sigma_{vis}}$$


Cross section seen by detector
(measured)

➤ σ_{vis} is determined in dedicated fills based on beam parameters

Luminosity calibration with van der Meer method

- ▶ beams are scanned across each other and luminosity recorded in luminometers [1],
- ▶ beams overlap width can be extracted $\Sigma_{x,y}$, to calculate the transverse luminous area.
- ▶ aimed to obtain the detector-specific visible cross-section
- ▶ rate can be correlated with instantaneous luminosity from beam parameters:

$$\sigma_{vis} = \frac{\mu_{pk}}{n_1 n_2} \times 2\pi \Sigma_x \Sigma_y \rightarrow \mathcal{L}_{inst} = \frac{\mu_{pk} f_{rev}}{\sigma_{vis}}$$




- ▶ beam-related systematic effects have to be considered.

Luminosity calibration with van der Meer method

- ▶ beams are scanned across each other and luminosity recorded in luminometers [1],
- ▶ beams overlap width can be extracted $\Sigma_{x,y}$, to calculate the transverse luminous area.
- ▶ aimed to obtain the detector-specific visible cross-section
- ▶ rate can be correlated with instantaneous luminosity from beam parameters:

$$\sigma_{vis} = \frac{\mu_{pk}}{n_1 n_2} \times 2\pi \Sigma_x \Sigma_y \rightarrow \mathcal{L}_{inst} = \frac{\mu_{pk} f_{rev}}{\sigma_{vis}}$$

- ▶ beam-related systematic effects have to be considered.



Motivation - Introduction

collaborative work of all LHC
experiments within the LLCMWG

- precision luminosity measurement requires a thorough understanding of beam systematics
- of particular importance: detailed studies for corrections and uncertainties related to the **Beam-Beam (BB) interaction**
 - BB optical distortion corrections underestimated in Run 1-2
 - BB angular deflection known, measured very well and calculated analytical [3b]
 - year-long studies to derive new model and strategy for systematic uncertainties, resulted in nice publication [3]
 - leading to the shift of the absolute integrated luminosity by $\sim -1\%$ [2] (compared to pre-2021)





in preliminary Run-2 [ATLAS results](#) $\sim 1.5\%$
correction with 0.2% uncertainty (!)



in legacy Run-2 [ATLAS results](#) $\sim 0.5\%$
correction with 0.3% uncertainty


Beam-beam interaction

- BB force : electromagnetic interactions of the two charged beams
- Change in orbit [3b]
- Change in optical properties [3]
- LHC specific vdM with multiple experiments in collision
- BB parameter describes the linearised force for small amplitude particles, separation introduces more complex effects



Beam-beam interaction

- BB force : electromagnetic interactions of the two charged beams
 - Change in orbit [3b]
 - Change in optical properties [3]
 - LHC specific vdM with multiple experiments in collision
- BB parameter describes the linearised force for small amplitude particles, separation introduces more complex effects
- COMBI [4] code used to model self-consistently the interactions to understand and quantify the bias to absolute luminosity measurements with multiple IPs
- Provide a set of corrections to be used in detectors luminosity analysis :
 - vdM analysis of absolute calibration of luminometers ($\xi < 0.01/\text{IP}$)
 - Luminometers non-linearities in high pile-up regime ($\xi = 0.01/\text{IP}$)




BB bias to luminosity break down for single IP:

Beam-beam force will modify the luminosity while scanning introducing different effects.

Studied separately in terms of:

- Optical effects including dynamic-beta, non linear effects and overlap changes (non-gaussianity and non-factorisation)
- Orbit deflection calculated from Bassetti-Erskine formula [5]
- In addition while one experiment is scanning the others acquire luminosity and introduce further BB effects:
 - Change in tunes
 - Amplitude dependent beta-beating
 - Phase advance dependency...





BB bias to luminosity break down for single IP:

Beam-beam force will modify the luminosity while scanning introducing different effects.

Studied separately in terms of:



- Optical effects including dynamic-beta, non linear effects and overlap changes (non-gaussianity and non-factorisation)
- Orbit deflection calculated from Bassetti-Erskine formula [5]
- In addition while one experiment is scanning the others acquire luminosity and introduce further BB effects:
 - Change in tunes
 - Amplitude dependent beta-beating
 - Phase advance dependency...



Measurements in VdM
CERN-ACC-NOTE-2013-0006
J. Wenninge, Kozanecki, Pieloni

Multi-collision study for vdM calibration

- focus on the additional collisions at interaction points (IPs) other than the scanning IP
- separate corrections for beam-separation dependent deflection-induced orbit shift and optical distortion (aka dynamic-beta)




Additional collisions \rightarrow IPs are coupled via BB

- additional betatron tune shift [6]
- Amplitude dependent beta-beating propagated
- Propagates from one IP to the others: phase advance between IPs causes modulation calibration constant [7]

Mimicking multi-IP impact

luminosity bias correction model based on the single-IP parametrization


- dependent on beams separation, BB parameter and tunes [3]
- effective multi-IP tune shift can be used to obtain the equivalent calibration constant bias (mimic the extra HO with a tune shift $0.5 \square / \text{NSIP}$)
- simple scaling law derived from strong-strong simulations
 - valid for all LHC IPs
 - verified in simulation for vdM regime $\xi \sim 0.004/\text{IP}$



If you cannot measure it, it doesn't exist!


Benchmark experiment

- Test designed especially to measure the BB effects
- phase advance between IP1 & IP5 optimised so as to **maximize** the effect on luminosity at the observer IP at injection energy
 - lattice validated (R. Tomas, T. Person, OP crew)



Multiple instruments were used to measure the BB effects on:


- luminosity from ATLAS and CMS luminometers
- tune spectra from ADT, BBQ
- transverse beam sizes with synch. light monitors and wire scanners
- orbit at the IPs with BPMs



W. Yi EPFL TPIV projects 2022

Benchmark experiment

- Test designed especially to measure the BB effects
 - phase advance between IP1 & IP5 optimised so as to **maximize** the effect on luminosity at the observer IP at injection energy
 - lattice validated (R. Tomas, T. Person, OP crew)



Measured beta-beating along the LHC ring from the knob

	Beam 1		Beam 2	
	$\Delta\mu_x [2\pi]$	$\Delta\mu_y [2\pi]$	$\Delta\mu_x [2\pi]$	$\Delta\mu_y [2\pi]$
IP1-IP5	30.977	29.649	31.062	29.762
IP1-IP5 adjusted	30.9	29.9	30.9	29.9
expected change	-0.077	0.251	-0.162	0.138
measured change	-0.076 ± 0.003	0.240 ± 0.002	-0.162 ± 0.002	0.137 ± 0.002




Figure 2: Measured beta difference between the lattice with the maximizing (+1) phase knob and nominal lattice along the LHC ring, for Beam 1 (left) and Beam 2 (right).

Multiple instruments were used to measure the BB effects on:

- luminosity from ATLAS and CMS luminometers
- tune spectra from ADT, BBQ
- transverse beam sizes with synch. light monitors and wire scanners
- orbit at the IPs with BPMs

Measured beta-beating along the LHC ring from the knob with reference to the MADX model predictions




Figure 5: Measured beta function differences along the LHC ring with respect to the MADX model with included maximizing (+1) phase knob, for Beam 1 (left) and Beam 2 (right).


Benchmark experiment

Series of tests:




- Scanning IP : in and out collision and transverse scan

→ propagation

- Witness IP: in HO collision, observation point to see bias on luminosity



Tune shift induced by BB during separation scan in horizontal plane at one IP, while the other is colliding head-on as measured by the ADT ObsBox[9]



Benchmark experiment


Series of tests:

- Scanning IP : in and out collision and transverse scan

→ propagation

- Witness IP: in HO collision, observation point to see bias on luminosity


Tune shift induced by BB during separation scan in horizontal plane at one IP, while the other is colliding head-on as measured by the ADT ObsBox[9]




Tune shifts and coherent modes

- Tune spectra and coherent modes
- Tune shift versus separation scan

COMBI simulated



Data




Benchmark experiment


witness	IP5				IP1				IP5			
ξ_{start}	0.010				0.007				0.006			
step #	2	4	6	8	2	4	6	8	2	4	6	8
bias [%]	3.21	3.58	3.01	3.41	2.42	2.64	2.34	1.98	2.46	1.89	1.37	2.23
stat. [%]	0.45	0.35	0.33	0.32	0.34	0.35	0.35	0.35	0.45	0.39	0.46	0.42

CMS luminosity change as a function of the ATLAS collision

$$\xi = 0.01/IP$$




$$\xi = 0.006/IP$$



ATLAS luminosity change as a function of the CMS collision

Luminosity observations $\xi = 0.007/IP$






- Luminosity bias due to BB has been observed in both observing IPS and the resulting effect is in within expectations
- The expectation varies with ξ_{bb}
- Phase advance impact to the observed effect visible

Benchmark experiment

presented at EPS-HEP 2023

Aim: validation of the correction strategy used in the vdM calibration

- support for the multi-IP modelling
- scaling law with BB parameter verified
- observations of BB-induced changes during a separation scan
- **first measurement** of the impact of BB effects on the luminosity in LHC





Extrapolation to nominal conditions



At nominal conditions the luminosity measurement can be biased with a non-linearity of a detector response over a wide pile-up range

- BB simulations useful to produce dedicated corrections - minimising the associated extra systematic from bunch by bunch differences
- Tested/used for a specific measurement fill (BSRT calibration fill 2023)

$$\sigma_{vis} = 2\pi \frac{\mu_{pk}}{n_1 n_2} \Sigma_x \Sigma_y$$



Impact of BB on detector non-linearity



Pile-up (PU) = ~ 7 x Single Bunch
Instantaneous Luminosity (SBIL)

Proof of concept (EPS-HEP 2023 J. Wanczyk)

- **apparent BB-induced slope** - removed with BB simulation predictions ($\xi \sim 0.008$)
 - fundamental to understand for HL-LHC
- Other luminometers behave differently

Independent measurement \rightarrow further studies needed for precise measurement

Conclusions

- Extensive simulation campaign of BB effects on the luminosity led to a much better understanding, minimising the related **systematic uncertainty** on absolute luminosity calibrations at LHC exp
- Improved corrections
 - optical effect shifted pre-2021 central values by -1% - improved results from ATLAS already published [2], CMS results on the way
 - by accounting for the multiple collisions effects - additional 0.4% correction for typical vdM BB parameter $\xi \sim 0.004/\text{IP}$
- Dedicated BB experiment at the LHC allowed to **validate some key aspects of the simulation model at the % level**
 - **First measurement** of the beam-beam-induced biases on luminosity
 - agreement with the simulation to the level of 0.1%
- Beam-beam simulations allow for dedicated corrections at the physics conditions (dedicated mini scan at $\xi \sim 0.01/\text{IP}$)
- Possible to remove the apparent beam-beam induced bias to detector response \rightarrow measuring intrinsic detector non-linear response in an independent way
 - luminometers non-linearities are expected to be one of the main challenges at HL-LHC
- Numerical simulations are invaluable tools to improve understanding, quantify effects and push higher precisions \rightarrow full exploitation of LHC luminosity and learn more in preparation for the high pile-up era
- BB induced Lumi enhancement by tuning the IPs can be applied also to LHC and HL-LHC case \rightarrow 3-7% depending on leveling at IPs

Thank you!

References

- [1] S. Van der Meer, “[Calibration of the Effective Beam Height in the ISR](#)” *CERN-ISR-PO-68-31*, 1968.
- [2] [ATLAS Run 2 luminosity calibration](#) / CMS on the way
- [3] A. Babaev et al., [arXiv:2306.10394](#), submitted to EPJC
- [3b] J. Wenninger, SL Note 96-01 (OP)
- [3b] M. Venturini and W. Kozanecki, SLAC-PUB-8700
- [4] T. Pieloni, [COMBI](#)
- [5] X. Buffat, 6D BB models:
- [6] W. Herr, [CAS proceedings](#)
- [7] J. Warczyk, [Phase modulation](#)
- [8] G. Trad, [BSRT](#)
- [9] M. Söderén et al., [ADT](#)